Trading Ratios: Salt Lake City PM_{2.5} Maintenance Plan

The Utah Division of Air Quality performed an air quality model (CAMx 6.30) analysis to estimate 24-hr $PM_{2.5}$ concentrations given reductions in on-road mobile NOx, VOCs, and direct $PM_{2.5}$ emissions. Reductions in on-road mobile emissions didn't include refueling or resuspended road dust.

Three different simulations were conducted using the 2035 Salt Lake City nonattainment area (SLC NAA) emissions inventory (EI). Each modeled simulation involved taking a 1 ton/day reduction in one of three on-road mobile EI pollutants: NOx, VOC, and direct PM_{2.5}. Resultant reductions in modeled PM_{2.5} were evaluated at the Rose Park FRM monitor, which is the monitor that consistently measures the highest value in the SLC NAA. Reductions were evaluated for the modeled episode day with the largest 24-hour PM_{2.5} concentrations at the Rose Park monitor. The reduction in modeled 24-hour PM_{2.5}, given a 1 ton/day reduction in a specific on-road mobile EI pollutant, is shown in Table 1:

El Pollutant	Modeled PM _{2.5} Reduction (µg/m³)*
NOx	0.152
VOC	0.046
Direct PM _{2.5}	0.955

Table 1: Relative contributions of NOx, VOC, and direct PM_{2.5} emission (1 ton/day) reductions to modeled PM2.5 concentrations at SLC NAA monitors.

To establish the MVEB ratios, simple division was applied between the model sensitivity results.

NOx : PM2.5 trading ratio = 0.955 / 0.152 = 6.3

VOC: PM2.5 trading ratio = 0.955 / 0.046 = 20.9

Future increases in on-road direct $PM_{2.5}$ emissions may be offset with future decreases in NO_x emissions from on-road mobile sources at a NO_x to $PM_{2.5}$ ratio of 6.3 to 1 and/or future decreases in VOC emissions from on-road mobile sources at a VOC to $PM_{2.5}$ ratio of 20.9 to 1.

^{*}These numbers have been rounded. The ratios were calculated using number that were not rounded